

Knollenbonitur der Öko-Kartoffeln im Sortenversuch nur Standort Willich (VIE) 2014

Sorte	Reife- gruppe*	Koch- typ**	Rizocto- nia %	Drycore %	Drahtwu- rm %	Schorf %	Schorf Index	Rhizoctonia Index	Drycore/ Drahtwurm Index
Alexandra	sf	f	50	15	0	90	1,7	1,1	1,0
Andrea	sf	f	73	10	1	33	1,2	1,2	1,0
Suzan	sf	f	85	7	4	7	1,0	1,0	1,0
Agila ¹	f	f	64	5	2	6	1,1	1,4	1,0
Fidelia	f	f	81	25	2	84	1,3	1,1	1,0
Goldmarie	f	f	92	5	2	5	1,1	1,0	1,0
Isabelia	f	f	48	6	2	1	1,0	1,4	1,0
Venezia	f	f	45	27	6	64	1,1	1,1	1,0
Christel	f	vf	80	11	5	50	1,2	1,1	1,0
Cardinia	f	vf	87	12	7	2	1,0	1,1	1,0
Queen Anne	f	vf	70	19	5	64	1,0	1,0	1,0
Wega	f	vf	81	19	2	23	1,5	1,1	1,0
Allians ¹	mf	f	63	13	7	99	2,3	1,2	1,0
Ditta ¹	mf	f	92	14	5	86	1,2	1,0	1,0
Ramona	mf	f	70	2	0	77	1,7	1,2	1,0
Torenia	mf	f	86	5	5	8	1,0	1,0	1,0
Belmonda	mf	vf	90	19	2	73	1,0	1,0	1,0
Birgit	mf	vf	65	18	7	86	2,1	1,3	1,1
Caprice	mf	vf	74	9	3	25	1,1	1,1	1,0
El Mundo	mf	vf	32	10	2	100	1,8	1,0	1,0
Georgina	mf	vf	78	13	1	0	1,0	1,1	1,0
Lilly	mf	vf	73	8	1	10	1,2	1,2	1,0
Salute	mf	vf	14	8	1	6	1,0	1,3	1,0
Mittel der Sorten			69	12	3,1	43,4	1,3	1,1	1,0

* sf = sehr früh, f = früh, mf = mittelfrüh, ms ** f = festkochend, vf = vorwiegend festkochend, m = mehligkochend